

American Academy of Child and Adolescent Psychiatry

AACAP is pleased to offer Practice Parameters as soon as they are approved by the AACAP Council, but prior to their publication in the *Journal of the American Academy of Child and Adolescent Psychiatry (JAACAP)*. This article may be revised during the *JAACAP* copyediting, author query, and proof reading processes. Any final changes in the document will be made at the time of print publication and will be reflected in the final electronic version of the Practice Parameter. AACAP and *JAACAP*, and its respective employees, are not responsible or liable for the use of any such inaccurate or misleading data, opinion, or information contained in this iteration of this Practice Parameter.

PRACTICE PARAMETER FOR CULTURAL COMPETENCE IN CHILD AND ADOLESCENT PSYCHIATRIC PRACTICE

ABSTRACT

The United States (U.S.) faces a rapidly changing demographic and cultural landscape, with its population becoming increasingly multi-racial and multi-cultural. Consequently, cultural and racial factors relating to mental illness and emotional disturbances deserve closer attention and consideration. In this Practice Parameter, we outline clinical applications of the principle of cultural competence that will enable child and adolescent mental health clinicians to better serve diverse children, adolescents, and their families. **Key words:** culture, ethnicity, race, cultural diversity, cultural competence, acculturation, ethnopharmacology.

ATTRIBUTION

This parameter was developed by Andrés J. Pumariega, M.D., Eugenio Rothe, M.D., Ayesha Mian, M.D., Lee Carlisle, M.D., Claudio Toppelberg, M.D., Toi Harris, M.D., Rama Rao Gogineni, M.D., Sala Webb, M.D., Jacqueline Smith, M.D., and the American Academy of Child and Adolescent Psychiatry (AACAP) Committee on Quality Issues (CQI): Heather Walter, M.D., M.P.H. and Oscar G. Bukstein, M.D., M.P.H., Co-Chairs; and Christopher Bellonci, M.D., R. Scott Benson, M.D., Regina Bussing, M.D., Allan Chrisman, M.D., John Hamilton, M.D., Munya Hayek, M.D., Helene Keable, M.D., Nicole Quiterio, M.D., Carol Rockhill, M.D., Ulrich Schoettle, M.D., Matthew Siegel, M.D., and Sandra Stock, M.D.

AACAP Practice Parameters are developed by the AACAP CQI in accordance with American Medical Association policy. Parameter development is an iterative process between the primary author(s), the CQI, topic experts, and representatives from multiple constituent groups, including the AACAP membership, relevant AACAP Committees, the AACAP Assembly of Regional Organizations, and the AACAP Council. Details of the parameter development process can be accessed on the AACAP website. Responsibility for parameter

content and review rests with the author(s), the CQI, the CQI Consensus Group, and the AACAP Council.

The AACAP develops both patient-oriented and clinician-oriented Practice Parameters. Patient-oriented parameters provide recommendations to guide clinicians toward best assessment and treatment practices. Recommendations are based on the critical appraisal of empirical evidence (when available) and clinical consensus (when not), and are graded according to the strength of the empirical and clinical support. Clinician-oriented parameters provide clinicians with the information (stated as principles) needed to develop practice-based skills. Although empirical evidence may be available to support certain principles, principles are primarily based on clinical consensus. This parameter is a clinician-oriented parameter.

The primary intended audience for the AACAP Practice Parameters is child and adolescent psychiatrists; however, the information contained therein may also be useful for other mental health clinicians.

The authors wish to acknowledge the following experts for their contributions to this parameter: Ian Canino, M.D., Terry Cross, MSW, Mario Hernandez, Ph.D., Roberto Lewis Fernandez, M.D., and John Sargent, M.D.

Jennifer Medicus served as the AACAP staff liaison for the CQI. This Practice Parameter was reviewed at the Member Forum at the AACAP Annual Meeting in October 2011.

From November 2012 to January 2013, this parameter was reviewed by a Consensus Group convened by the CQI. Consensus Group members and their constituent groups were as follows: Heather J. Walter, M.D., M.P.H., chair, Christopher Bellonci, M.D., Carol Rockhill, M.D., Helene Keable, M.D. (CQI); Terry Cross, John Sargent, M.D. (Topic Experts); Kathlene Trello-Rishel, M.D., Gabrielle Shapiro, M.D. (AACAP Assembly of Regional Organizations); Steven Adelsheim, M.D., Warren Y. K. Ng, M.D. (AACAP Council).

Disclosures of potential conflicts of interest for authors and CQI chairs are provided at the end of the parameter. Disclosures of potential conflicts of interest for all other individuals named above are provided on the AACAP website on the Practice Information page.

This Practice Parameter was approved by the AACAP Council on April 16, 2013.

This Practice Parameter is available on the internet (www.aacap.org).

Correspondence to the AACAP Communications Department, 3615 Wisconsin Avenue, NW, Washington, D.C. 20016.

©2013 by the American Academy of Child and Adolescent Psychiatry.

INTRODUCTION

The rapidly changing demographics of the U.S. are largely a result of three major factors: progressive aging and low birth rate of the European-origin population, lower mean ages and increasing birth rates in non-European minority groups, and a significant rise in immigration from non-European countries, especially from Latin America, Asia, and Africa. By 2050, European-Americans will no longer constitute the majority, and this will happen by 2030 among children under eighteen and is already true among children under eight.¹ The process of evaluating and treating culturally diverse children and youth and their families can be complex and requires special expertise and unique approaches. Thus, this parameter can be useful for clinicians and, ultimately, for the children and families they serve. Principles in this parameter apply to culturally diverse children and youth under the age of 18.

METHODOLOGY

In PubMed the MeSH terms culture, Hispanic, Latino, African American, Asian American, American Indian, child psychiatry, child psychology, adolescent psychiatry, adolescent psychology, and United States were searched. The initial search yielded 2,970 results. Results were then limited to English, human, “all child (0 to 18 years)” and 1990 through December 2011. Additional limits included: classical article, clinical trial, comparative study, controlled clinical trial, evaluation studies, guideline, historical article, meta analysis, practice guideline, multicenter study, randomized controlled trial, review, twin study and validation studies. The refined PubMed search yielded 2,268 articles.

In the PsycInfo database subject headings (focused) the keywords "culture", "Latino", "Hispanic", "African American", "Asian American", "American Indian", and "Mental Health" were searched. The initial search returned 40,167 articles and was then limited to English, articles in the United States, “childhood: birth to age 12 yrs,” “adolescence: age 13-17 yrs,” “peer reviewed journal,” and 1990 through December 2011. The refined PsycInfo search yielded 2,240 articles.

In the Cochrane Database of Systematic Reviews, keywords of culture and mental health were searched without additional limits. The Cochrane search yielded 80 articles. An additional 953 articles were retrieved from the CINAHL database, after excluding Medline articles, by searching “culture”, “Latino”, “Hispanic”, African American", "Asian American", "American Indian", "Mental Health" , and "United States", limited to childhood and adolescence and limiting to peer reviewed articles, English language, and 1990 through December 2011.

A total of 5,461 articles were identified. After removing duplicate references the resulting yield from the comprehensive search was 4,391 articles. The titles and abstracts of all articles were reviewed. Studies were selected for full text review based upon their place in the hierarchy of evidence (e.g., randomized controlled trials), quality of individual studies, and generalizability to clinical practice. The search was augmented by review of articles nominated by expert reviewers and further search of article reference lists and relevant textbook chapters. A total of 163 articles were selected for full text examination. Principles were identified from the consensus of the AACAP Diversity and Culture Committee, informed by the literature review articles and the Cultural Competence Standards for the Four Racial/Ethnic Underserved Underrepresented Racial/Ethnic Groups.²

DEFINITIONS

- Culture: Integrated pattern of human behaviors including thoughts, communication, actions, customs, beliefs, values, and institutions of a racial, ethnic, religious, or social nature.
- Cultural competence: Set of congruent behaviors, attitudes, and policies found in a system, agency, or professionals that enables them to work effectively in a context of cultural difference.³
- Acculturation: Process of change in the cultures of two or more groups of individuals from different cultures, resulting from their continuous first-hand contact.⁴
- Race: Social classification system based on a set of external physical characteristics that are socially significant within a specific culture.
- Ethnicity: Common historical or geographic heritage shared by a group of people.
- Immigrant: Someone who intends to reside permanently in a new land.

HISTORICAL OVERVIEW

The recent demographic changes in the U.S. are highly significant for child mental health services. First, the acceptability and use of mental health services are highly governed by cultural attitudes, beliefs, and practices. Second, the current science base around psychiatric diagnosis and treatment is derived from research primarily involving European-origin populations, so its validity for these emerging populations is not fully established. Lastly, minority populations face many increasing challenges around mental illness, including different sources of stressors, changing patterns of psychopathology, lower access to services and evidence-based treatments, and higher burdens of morbidity and possibly mortality than Euro-Americans. For example, Latino and African American youth have significantly higher rates of suicidal ideation and attempts compared to Euro-Americans.⁵

The current health care system does not effectively address the needs of culturally diverse populations. The recognition of racial and ethnic disparities in general health care has led to increasing recognition of similar disparities in mental health care.^{6,7} One obvious example of mental health disparities are the child welfare and juvenile justice systems. Over 50% of children and youth in the child welfare system are African-American, Latino, and American Indian children, and over 65% of the children and youth in the juvenile justice system are African-American and Latino.^{1,8} These systems also serve disproportionate numbers of mentally ill children, and their racial ethnic disparities are associated with a lack of early access to mental health services to prevent such outcomes. At the same time, in response to these mounting challenges, *cultural competence* became one of the core principles of the children's community-based systems of care movement.³

PRINCIPLES

Principle 1. *Clinicians should identify and address barriers (economic, geographic, insurance, cultural beliefs, stigma, etc.) that may prevent culturally diverse children and their families from obtaining mental health services.*

Non-Hispanic white families are twice as likely as minority families to seek mental health treatment for their child⁹⁻¹¹ despite evidence suggesting the prevalence of psychiatric disorders in children does not appear to vary greatly by race or ethnicity.¹² When minorities seek treatment, they may not remain engaged in outpatient services or utilize as many service units.^{10,11} Multiple systemic and logistical barriers that interfere with timely access to services are disproportionately experienced by racially/ethnically diverse families. These include

financial needs, location of services and transportation, lack of adequate insurance, poorly understood bureaucratic procedures, and lack of linguistic support. Even when minority children and families receive services, these are often interrupted prematurely by these barriers.¹³

Though many traditional cultural values and beliefs are a source of strength and support for diverse children and families, some can act as barriers to mental health services. For example, a survey of Latinos found factors such as perceptions of mental illness (including stigma and beliefs about causality), fatalism, spirituality, familism (in which the family is considered the primary unit of identification and allegiance³ and leads to keeping problems within the family), cultural commitment (e.g., to using only culturally sanctioned helping approaches), and language proficiency¹⁴ affect their treatment. Abe-Kim et al.¹⁵ found that rates of mental health–related service use, subjective satisfaction, and perceived helpfulness of services varied by birthplace and generation; US-born Asian Americans used services at higher rates than their immigrant counterparts, and third-generation or later individuals had the highest (62.6%) rates of service use in the previous year.

Stigma can be a powerful barrier to timely access to treatment. In many cultures, mental illness has major negative connotations, leading to the fear of double-discrimination (as a result of being both culturally different and perceived as “crazy”), which prevents minority families from accessing services. These perceptions can then become self-reinforcing when emergency services are needed due to the traumatic impact of suddenly receiving a more restrictive level of care and/or their involuntary nature.¹⁶ Families may mistrust mental health service agencies given their past histories of discrimination and disregard for cultural needs.¹⁷ Culturally diverse families are more vulnerable to perceived or actual power differentials in their encounters with health care professionals.

Diverse families should be educated to improve their understanding of diagnosis and treatment, empower decision-making, expose myths, and improve treatment outcomes.² Clinicians should address the realities and perceptions of power differentials that may interfere with therapeutic relationships.⁸

It is not uncommon for minority children and adolescents to engage in mental health treatment for troubling behaviors identified by the school or the court, rather than by their parents.¹³ One explanation is that minority families tend to have a higher threshold for disruptive behaviors and to not seek professional intervention until the situation becomes unmanageable.¹⁸

Yeh et al.¹⁹ reported that children and adolescents of ethnic and racial minorities are referred for mental health treatment from involuntary sources such as the child welfare and juvenile justice systems more often than their white counterparts, with rates of self-referral to community services that are lower than those for whites. Though socioeconomic class can contribute to these disparities, they affect minority youth at higher rates than white youth even controlling for socioeconomic status.

Principle 2. Clinicians should conduct the evaluation in the language in which the child and family are proficient.

Language-based communication is critical in obtaining accurate clinical information and establishing a therapeutic alliance. Many immigrants, however, are not fluent in English and, thus may not be able to fully participate in the clinical process. In these situations, translation and interpretation are critical to effective care. Limited English proficiency is a significant barrier to accessing mental health services for adults and children from different ethnic origins, resulting in significantly lower utilization of mental health services.^{11,14,20} Lack of appropriate linguistic ability or interpreter support has been associated with misdiagnosis and adverse clinical outcomes.^{21,22} In these cases, clinicians should either obtain linguistic support through qualified interpreters, or possess demonstrable proficiency in the target language. Unfortunately, translation and interpretation are oftentimes considered menial or informal tasks in the clinical process, as reflected by using untrained interpreters or translators without regard to impact on family relations, family members, siblings, or the child.² Language brokering, the common practice of having children act as interpreters between parents and medical and school authorities, should be avoided, particularly when the patient is the language broker. An association has been identified between high language brokering contexts and higher levels of family stress, lower parenting effectiveness, poorer adjustment in academic functioning, higher Child Behavior Checklist (CBCL) internalizing scores, and substance use in adolescents.²³ Telephonic interpretation services enable 24 hour access and a wide range of available languages, but are not ideal due to their lack of ability to convey nonverbal communication. Interpreters should have proper training in both the skill of interpretation and the content area being discussed. They should serve as integral members of the clinical team, serve as cultural consultants when they have understanding of the family's culture, and interpret all of the verbal,

non-verbal, and implicit communications from the child and family rather than provide summaries.²⁴ If live translation services are not available, clinicians may have to use other alternatives (telephonic interpreter services, individuals with dual language abilities, or someone the family or child identifies), but efforts should be made to obtain the consent of the individuals using written or non-verbal means. Any educational materials and rating or diagnostic instruments should be translated to the language of the family member or child, and their reliability and validity in the target cultural group should be established using well accepted reverse translation and psychometric methodologies. In the event materials or instruments have not been formally translated or validated, practitioners should access translation services from trained professionals or from bilingual child mental health professionals. Clinicians should be cautious about clinical interpretations based on diagnostic instruments not properly translated or validated with the population in question.^{2,24}

Principle 3. Clinicians should understand the impact of dual language competence on the child's adaptation and functioning.

An estimated 20% of American children 18 and under grow up exposed to two languages. Learners of English as a “second language” (defined as any language learned after age 3, which is the end of the critical period for rapid language acquisition) constitute the majority of dual language children in America. After English, the most common home language in the United States is Spanish.^{25,26} Grammatical and other language errors made by a child learning a second language (or a second English dialect, such as standard American English for a speaker of Black English) should not be confused with the grammatical or lexical abnormalities of language disorders. On the other hand, deficits associated with psychiatric and language disorders (such as auditory-verbal working memory deficits) may slow the acquisition of a second language. Specialized consultation and assessment over time by a speech/language pathologist with expertise in dual language children may be necessary to differentiate normal from disordered language acquisition.²⁶

There is evidence that maintaining the first (home) language is important in accessing family and community protective factors and other benefits. Despite this evidence, there has been a poorly substantiated practice of recommending to parents that they discontinue speaking the home language to a child who is facing language, cognitive or other delays. This practice has

little or no empirical support, and the limited research conducted in this area suggests that children with language impairment can be healthily exposed to and learn two languages with no significant detrimental effects.²⁶ While it may be true that certain children with linguistic or other deficits may become overwhelmed by the additional cognitive and linguistic demands of dual language learning, recommendations to discontinue learning the home language may have potentially serious consequences and should not be made lightly. Rather, such decisions should ideally involve full assessment by a speech/language pathologist with appropriate expertise, consultation with the parents and others who know the child well, and an informed decision process by the parents with consideration to the family's plans for the future.²⁶ For example, a child whose family only speaks a minority language may need to maintain and learn that language.

Sudden language immersion may be stressful, depending on the child's temperament and availability of supports. Children who are suddenly immersed in a second language environment with no knowledge of the language may go through a normal "nonverbal period." This should not be confused with selective mutism, which has a higher prevalence among immigrant dual language children. Thus, it is important that clinicians be familiar with features that differentiate the normal nonverbal period from selective mutism, which typically lasts much longer.²⁷

Principle 4. Clinicians should be cognizant that cultural biases might interfere with their clinical judgment, and work towards addressing these biases.

Both adult and pediatric psychiatry literatures provide evidence for cultural and racial disparities in diagnostic assessment, treatment measures, and the quality of received health care.^{8,10,28} Stereotyping, biases and uncertainties in health care providers can lead to unequal treatment.^{6,29} When the patient's presentation or diagnosis is unclear, physicians may inadvertently over-rely on behavioral or clinical stereotypes of specific groups at the expense of focusing on the patient's unique experience, clinical presentation or socio-cultural context.

Stereotyping is defined as the process by which people use social categories (e.g., race, sex) to acquire, process, and retrieve information about others. Because stereotyping can be a subtle cognitive phenomenon resulting from virtually universal social categorization processes, it also occurs, often unconsciously, among people who strongly endorse egalitarian principles and truly believe that they are not prejudiced.³⁰ In the U.S., there is considerable empirical evidence

that well-meaning people who are not overtly biased and do not believe they are prejudiced, typically demonstrate implicit (i.e., unconscious) negative racial attitudes and stereotypes.³¹

Prejudice is defined in psychology as an unjustified negative attitude based on a person's group membership.³¹ While in general medicine evidence exists that provider biases adversely impact quality of care for minority patients,⁶ this has not been extensively studied in mental health. Research does suggest that behavioral and general healthcare providers' diagnostic and treatment decisions, as well as their feelings about patients, are influenced by race or ethnicity. For example, mental health professionals subliminally "primed" with African American stereotype-laden words were more likely to evaluate the same hypothetical patient (whose race was not identified) more negatively than when primed with neutral words.³²

Principle 5. Clinicians should apply knowledge of cultural differences in developmental progression, idiomatic expressions of distress, and symptomatic presentation for different disorders to the clinical formulation and diagnosis.

Motor, language, cognitive, and social development appear to differ across cultures. These differences may result from developmental expectations consistent with differentiated cultural value systems. Such differences, seen out of cultural context, could lead to the misidentification of pathology or misdiagnosis of mental disorders.^{24,33} Expressions for psychological or emotional distress also differ across cultures. Idioms of distress are linguistic or somatic patterns of experiencing and expressing illness, affliction, or general stress.³⁴ Idioms of distress do not generally correspond to diagnostic categories. They can at times be an alternative means of expression for disorders recognized by Western psychiatry, such as depression or anxiety disorders. However, they can just as easily be expressions of psychological distress unique to a given culture, or even common normative expressions of stress. More complex expressions of illness or distress are termed cultural syndromes.³⁴ Idioms of distress and cultural syndromes can often be mistaken for more serious psychopathology while they are often normal variants of expression of distress. For example, *ataques de nervios* (a reaction combining anxiety, agitation, and dissociation common amongst Latinos of Caribbean origin) can be confused with a psychotic reaction, while *falling-out* (an expression of emotional stress amongst Afro-Caribbeans that includes sudden acute paralysis and dissociation) can be confused with catatonia.

Common patterns of symptomatology displayed by children from diverse populations can differ significantly from those in Euro-Americans, partly due to the influence of their culture. For example, in minority youth, somatization and anger are frequently associated with depression and anxiety, and psychosis is often over-diagnosed.^{28,35} Emotional reactivity during depression is greater in Asian-origin individuals than in whites.³⁶

In the evaluation of children from less assimilated families it is critical to recognize cultural manifestations of distress or illness, such as cultural syndromes. Accurate assessment can prevent misdiagnosis and erroneous treatment. It is important to seek consultation from a colleague with cultural expertise on the cultural context and significance of particular idiomatic expressions or symptoms that may be culturally influenced.²

The DSM Outline for Cultural Formulation provided a useful framework for assessing sociocultural aspects of the patient's mental health and relating these to the clinical formulation^{37, 38}. The DSM 5 provides a revised version³⁹ that calls for systematic assessment of five distinct categories: 1) *cultural identity of the individual*, comprising the individual's racial, ethnic, or cultural reference groups as well as other relevant aspects of identity, such as degree of involvement with the culture of origin versus host culture, religion, socioeconomic background, place of origin, migrant background, and sexual orientation; 2) *cultural conceptualization of illness*, including the influence of cultural beliefs on the individual's experience, conceptualization, and expression of symptoms or problems; 3) *psychosocial stressors and cultural features of vulnerability and resilience*, including key stressors and supports in the individual's socio-cultural environment (such as religion, family, and social supports); 4) *cultural features of the relationship between the individual and the clinician*, such as dynamics of differences based on cultural, socio-economic, language, and social status that may cause differences in communication and influence diagnosis and treatment; and 5) *overall cultural assessment* summarizing the implications of the above aspects for diagnosis, plan of care, and other clinically relevant issues.

DSM 5 also includes the Cultural Formulation Interview (CFI),³⁹ a 16 item semi-structured interview to obtain information about the impact of culture on clinical presentation and care, with suggested questions to inquire about different domains. The CFI contains four assessment domains: a) cultural definition of the problem; b) cultural perceptions of cause, context, and support; c) cultural factors affecting self-coping and past help-seeking; and d)

cultural factors affecting current help-seeking. The CFI also includes Supplementary Modules as adjuncts to the core CFI that provide more in-depth assessment of various areas. The CFI was field-tested for diagnostic utility among clinicians and acceptability amongst patients as part of the DSM field trials, though the trials did not include children and youth.

An important reference for cultural differences in developmental progression, expressions of distress, and symptomatic presentation by gay, lesbian, and transgendered youth, and how these impact their assessment and treatment, can be found in the AACAP Practice Parameter for gay, lesbian, and transgendered youth.⁴⁰

Principle 6. Clinicians should assess for a history of immigration-related loss or trauma and community trauma (violence, abuse) in the child and family, and address these concerns in treatment.

Migration has risks and stressors unique to immigrants. These can be divided into: 1) pre-migration stress, such as exposure to violence, persecution and torture in the country of origin; 2) migration stress, including the disruption and separation of families, traumatic journeys, detention in refugee camps and various forms of victimization, such as abuse by smugglers; 3) post-migration and acculturation stress, resulting from low levels of education and job skills,⁴¹ living in high risk neighborhoods, with high exposure to crime and violence, and overcrowded, poor quality inner city schools.⁴² Additionally, immigrant children and their families often face the stressors of prejudice and discrimination against immigrants as well as those experienced by non-immigrant minorities, especially if racially different from the majority culture,⁴³⁻⁴⁵ which may lead to poor academic functioning, low self-esteem, depression and suicide.⁴⁴ Some minority youth adopt a position of defiance against the dominant culture by joining gangs.⁴⁶ As a result of these stressors, second-generation children of immigrants are generally at increased risk for mental health problems, including anxiety, depression, substance abuse, and posttraumatic stress disorder (PTSD).⁴⁷ The stresses of immigration on the family can also result in domestic violence and criminal activity.⁴⁸ Factors such as parental emotional well-being and peer relationships can mitigate or aggravate these problems.⁴⁴

Immigration to the U.S. can lead to complex patterns of family fragmentation and reunification. To get established, parents may immigrate before their children, leaving them with relatives in the country of origin. Children are often brought to the U.S. years later with hopes of

offering them a better education and future. Children are often torn away from grandparents and other extended family members who have become their primary caregivers, often with little recognition of the mourning process involved. Parents can become dismayed and disappointed with the unhappiness, even rejection, their estranged children may demonstrate towards them.⁴⁹ Children who have experienced such separations or losses have a higher risk for later developing depression, conduct, and substance abuse disorders compared to other immigrant children.^{44,49} Adolescent victims of war and genocide have elevated rates of conduct disorder, aggressive and sexual acting out behaviors, substance abuse, depression and PTSD.^{50,51} Those exposed to traumatic crossings and confinement in refugee camps experienced silent symptoms of PTSD, which often went unnoticed by teachers.^{43,50,51} Treatment of immigrant and refugee children as well as inner city minority children must address traumatic exposure from these various sources, use evidence-based interventions that address the impact of trauma in the child and family, and also provide for community support.⁵¹

Principle 7. Clinicians should evaluate and address in treatment the acculturation level and presence of acculturation stress and intergenerational acculturation family conflict in diverse children and families.

Acculturation directly impacts the developmental task of identity, just as racial/ethnic identity comprises a significant aspect of psychological identity. Peers and family members serve as “mirrors” against which the self is reflected.^{4,44} For racially/ethnically diverse children and youth, this mirroring comes from two sources: the traditional cultural environment of the home and the mainstream cultural environment of peers, school, and the broader community, which can result in conflicting images. Diverse children and youth often face significant pressures to assimilate into mainstream society through media images and implicit threats of social and economic marginalization. In the process of acculturation, research supports the best adaptational outcomes for youth are associated with the development of a *bicultural identity*, where the immigrant youth remain rooted in their culture of origin (often mediated by learning the home language) but have the necessary knowledge and interpersonal skills to successfully navigate mainstream culture (i.e., a hyphenated identity).^{4,44} However, immigrant children and families face other potential outcomes: 1) *marginalization*, with separation from both cultures and some degree of ethnic identity diffusion; 2) *assimilation*, where the culture of origin is rejected and the

host culture is largely adopted, with the loss of language and customs of origin; 3) *separation*, where the host culture is rejected and there is a strong adherence to the traditional culture and language, often remaining bound to the “ethnic enclave”; and 4) *negative identification*, a form of marginalization where the youth is openly hostile to both cultures and adopts values and practices antithetical to both (e.g., adoption of gang culture).^{4,52}

Acculturation stress is the distress and internal conflict resulting from adaptation to a new host culture, including internal cultural value conflicts and external pressures to assimilate, as well as facing the host society’s hostility in the form of racism and discrimination. It can also result from the loss of protective aspects of the traditional culture that help with developmental stresses (e.g., religious prohibition against suicide) while not yet adopting those from the host culture.⁴ Acculturation stress can increase risk for psychopathology including substance abuse, depression, anxiety, suicidality, conduct disturbance, and PTSD.^{4,53} It can also aggravate pre-existing conditions. For example, the rates of PTSD decreased more in Bosnian than Southeast Asian immigrant youth who were victims of ethnic cleansing one year after immigration. Becker and colleagues⁵⁴ speculated that fewer cultural differences between the country of origin and the host country accounted for the better adaptation of Bosnian refugees to the U.S. A number of studies found associations between a stronger ethnic identity with lower acculturative stress, better academic performance, better psychological adaptation, and fewer externalizing and internalizing symptoms, but greater awareness of discrimination. Parental stress and family separations can adversely affect acculturative stress, while positive family relations can ameliorate it.^{4,44}

Many immigrant families are headed by parents with traditional cultural beliefs, who are experiencing the process of acculturation themselves but without their children's cognitive, linguistic, or emotional flexibility. Differences in acculturation between parents and children can lead to inter-generational acculturation conflict, which may result in psychological distress. Such acculturation discrepancies are most prominent during adolescence. One factor that may contribute to acculturation conflicts is assumption of a parentified role by a more assimilated immigrant child as a means of assisting their less assimilated parents to navigate the new cultural milieu of language, rules, and norms. Though parentification can be adaptive, it can represent an inversion in family roles and authority if maintained over time and contribute to heightened inter-generational conflict when parents attempt to assert their authority.^{44,55}

Differential acculturation between immigrant parents and their children leads to incongruence of values and expectations, resulting in intergenerational conflict and distancing. In some Asian-American families, differential acculturation is a predictor of negative mental health consequences for youth, such as school difficulties, gang involvement, depression, and suicidality. In Hispanic families, orientation to mainstream American culture, when coupled with the retention of Hispanic cultural practices and heritage, is associated with positive adjustment outcomes in youth, such as the absence of conduct disorders and substance abuse. Increased parental involvement, positive parenting, promoting biculturalism in Hispanic adolescents and their families, and reframing intergenerational difficulties as culturally based disagreements, has been efficacious in improving family functioning and adolescent behavior problems.⁵⁵⁻⁵⁷

Acculturation stress, inter-generational acculturation conflict, and interventions designed to address them should be routine in the assessment and treatment of children of immigrants or minority families. Interventions aimed at enhancing minority adolescents' mental health should focus on the intergenerational relationship, as it is a predictor of the youths' outcome. Family-based intervention, where intergenerational and intercultural conflict is explored and the value of communication within cultural expectations is underscored, is effective when working with adolescents.^{55,56}

Principle 8. Clinicians should make special efforts to include family members and key members of traditional extended families, such as grandparents or other elders, in assessment, treatment planning, and treatment.

In collectivist cultures, in which people view themselves primarily as members of groups and usually consider the needs of the group over the needs of individuals, extended family involvement may be the only acceptable model of addressing emotional and mental health problems. In contrast, in individualistic cultures, these problems are addressed by the individual or immediate relatives. Therapeutic engagement of diverse families may call for strategies not utilized in individualistic cultures. Involvement of extended family members and non-blood relatives with an equivalent emotional bond (i.e., 'fictive kin') is important in obtaining necessary collateral input for appropriate diagnosis, subsequent treatment recommendations, and treatment adherence.⁵⁸ Although clearly beneficial, this extended involvement may raise challenging issues of confidentiality. Confidentiality should be maintained in such a manner that

it does not interfere with communication and collaboration. Clinicians should explain the importance of confidentiality in psychotherapy to the family, so they are not perceived as "driving a wedge" between the child and the family, nor used by the patient to resist dealing with family issues. However, family conflict often predicts the use of mental health services for diverse patients, so it is important that clinicians ascertain whom the patient considers family and determine if they are a source of stress or support.⁵⁹

The mental health literature has explored the potential impact of culture on boundary-keeping practices, i.e., the maintenance of parameters that individuals or societies use to define interpersonal and professional boundaries. People from collectivist backgrounds view concepts that define such boundaries, such as time, confidentiality, dual-relationships, and self-disclosure, differently than those from individualistic cultures. These differences can affect the therapeutic process.⁶⁰ It is recommended that clinicians carefully negotiate boundaries and treatment parameters with culturally diverse patients and families.⁶¹ During negotiation, the providers should respect culturally established means of communication and family role functioning, but also foster family flexibility in dealing with their bicultural offspring around such issues. This process establishes a collaborative process involving the mental health provider and the family. Savin and Martinez⁶¹ recommend a 'graded-risk model' to address boundary issues when treating culturally diverse families. They encourage clinicians to evaluate potential crossings of professional or interpersonal boundaries, consider potential harm/benefit to the patient/professional relationship, and identify the presence/absence of exploitative or coercive components and professional intent and aspirations, all within the context of the boundary crossing. By utilizing this process model, clinicians can analyze whether a boundary crossing may be harmful or beneficial for treatment.

Accurate, valid consent forms translated into the patient's and family's primary languages as well as interpreter services are essential (see Principle 2). It may be important to allow the family extra time to consult traditional and family decision-makers in their consent process. These decision-makers have major influence in the family consent process. Actively avoiding their input can result in significant problems with treatment adherence. An overly paternalistic style on the clinician's part may lead a patient to not return or politely agree to the plan but then not adhere. The use of family psychoeducation is critical to addressing questions and culturally based myths and misconceptions about psychiatric medications, including the

rationale for their use, mechanism of action, and benefits versus risks for the child.^{2,62} Diverse families often have a background of limited treatment options and decision-making opportunities. Thus, it is especially important to assist them in developing knowledge and skills to address their child's mental illness, question perceived authority, and exercising their rights for second opinions or alternative treatment resources.^{17,44}

Principle 9. *Clinicians should evaluate and incorporate cultural strengths (including values, beliefs and attitudes) in their treatment interventions to enhance the child's and family's participation in treatment and its effectiveness.*

Clinicians should assume a posture of genuine cultural curiosity in learning about the family's traditional cultural beliefs/values. A culturally competent approach^{3,63} promotes the incorporation of beliefs, values, attitudes, and cultural rituals and practices within mental health treatment, using psychoeducational approaches to bridge traditional understanding of illness and its treatment, with Western conceptualizations and its medical/scientific model. The process of participatory or shared decision-making, where cultural aspects are integrated into the implementation of evidence-based practice, is crucial to enhance culturally competent care that ensures treatment engagement.⁶⁴ Many culturally competent psychotherapeutic approaches and interventions have been developed, including storytelling and cultural themes,^{65,66} symbolic rites of passage,^{67,68} and matching therapists according to their racial/ethnic background.

Culturally adapted treatment modalities may have some improved effectiveness compared to standard interventions, though the evidence is mixed. Two studies using a correlational approach indicate that ethnic match between client and therapist was associated with positive outcomes following youth- and family-based treatment.^{69,70} A meta-analysis⁷¹ of 76 studies evaluating the benefit of culturally adapted mental health interventions found a moderate benefit for culturally adapted interventions. However, two smaller meta-analyses⁷² comparing culturally adapted to standard interventions did not find significant differences in effect sizes.

Diverse cultural groups' explanatory models for mental health and illness can vary, invoking spiritual, supernatural, sociological, and interpersonal explanatory models. Such explanatory models often lead families to seek help for their children's problems from a spiritual healer, church elder, community leader, or relative, rather than from mental health professionals. Reasons for such preferences include greater acceptability of the healer's explanatory model of

illness, greater family support, less stigma for seeking services, and perceived greater rapidity and effectiveness. This preference is seen particularly among first generation immigrants and refugees, including Latinos, Asian-origin, American Indian, and some European-origin groups.¹³

Clinicians should consider consulting and collaborating with traditional healers (such as curanderos, santeros, or shamans), and including rituals and ceremonies in psychotherapy with children from more traditional backgrounds. Collaboration with indigenous traditional healers can ameliorate cultural loyalty conflicts within families and children, and also improve access to care in populations unfamiliar with or even mistrusting of the medical/psychiatric model. This is typically feasible when traditional healing methods complement or enhance (and do not directly conflict with) the effectiveness of Western psychotherapeutic and pharmacological interventions. Traditional healers are often reticent to identify themselves as collaborating with Western-trained clinicians. However, mutual respect and education in exchanging information and perspectives can foster collaboration.¹³

Principle 10. Clinicians should treat culturally diverse children and their families in familiar settings within their communities whenever possible.

Diverse children and families often prefer ethnically specific community clinics or clinics located within schools or ethnic neighborhoods. For example, a study of referral patterns in four Asian American groups demonstrated a clear preference for ethnic-specific mental health services.⁷³ School-based services are also generally well accepted and highly effective. School-based disaster mental health services after Hurricane Katrina that were culturally adapted and evidence-based were effective at reaching minority youth who otherwise might have not gained from these treatments.⁷⁴ Similar success was identified in the Cognitive Behavioral Intervention for Trauma in Schools (CBITS) program in diverse ethnic populations within inner city Los Angeles.⁷⁵

Clinicians should favor home- or community- based alternatives to hospitalization for minority children due to the integral nature of community and family to their identity and emotional supports. Any out of home placement or hospitalization should ideally be accomplished with family and youth cooperation. The principles of treatment in the least restrictive environment are especially important given how some treatments may remove youth from their community, extended family and traditional cultural supports.^{24,76} Involuntary

hospitalization should be avoided as it tends to re-exacerbate past traumas and contribute to new ones, as well as re-awakening historic mistrust of the mental health system.¹⁷ Psychiatrically hospitalized African-American youth have a significantly higher tendency to be admitted involuntarily, which enhance such fears and mistrust.²⁸

The cultural competence model³ and the principles of community systems of care⁷⁶ promote the use of community resources and cultural strengths to facilitate effective interventions with diverse children and youth, reinforcing adaptive cultural values and beliefs. Systems of care programs have some demonstrated effectiveness in improving access to care and improving outcomes for diverse children and youth. A meta-analysis of services utilization data versus home community composition for the federal Comprehensive Children's Mental Health Initiative communities⁷⁷ found that the program successfully reaches disadvantaged and minority youth and can bring substantial infrastructure to address youth mental health disparities. Systems of care communities serving Latino youth have also demonstrated equivalent clinical and functional outcomes as for white youth.⁷⁸

Principle 11. Clinicians should support parents to develop appropriate behavioral management skills consonant with their cultural values and beliefs.

Culturally diverse parents tend to accept and adhere more to behavioral management skills when these are consonant with their culture. For example, a randomized test of a culturally-adapted parent management intervention with Spanish-speaking Latino parents of middle school-aged youth at risk for disruptive behavior disorders, with half in the culturally adapted program and half in the control condition, revealed parents in the culturally adapted program reported higher levels of satisfaction and better outcomes in parenting skills and youth externalizing behaviors. Attendance was high, with 70% of families attending at least 80% of sessions.⁷⁹ Kumpfer et al.⁸⁰ showed that recruitment and retention was 41% higher in culturally-adapted family intervention programs compared to non-adapted versions.

Clinicians should advise families about parenting approaches that may be incongruent with Western culture, especially ones that are illegal. For example, the parenting attitudes of Asian Indian mothers living in the U.S. differ from those in India and found the latter favored corporal punishment, had more inappropriate expectations developmentally, and tended to reverse roles with their children (more frequently endorsing such items as “young children

should be expected to comfort the mother when she is feeling blue”).⁸¹ In some Muslim American families arranged marriages, the significance of showing respect to parents, and polygamy are issues that can cause conflicts.⁸² It is also important to address parent-child boundary issues that may become problematic with the child’s increasing acculturation. For example, as part of their emphasis on the family, Chinese immigrant parents sleep in closer proximity to their children, and place a greater emphasis on the inhibition of expression, less emphasis on children expressing their needs, and less importance on children’s intimacy and pleasure.⁸³

Principle 12. Clinicians should preferentially use evidence-based psychological and pharmacological interventions specific for the ethnic/racial population of the child and family they are serving.

The Surgeon General’s Supplement on Culture, Race, and Ethnicity⁷ pointed out the racial/ethnic disparity in the evidence base for various psychiatric and psychological interventions as a result of a lack of evaluation of such interventions in minority populations. Much of the research disparity has been the result of over-representation of Euro-Americans and inadequate recruitment of minorities into clinical trials, such that most of the presumed evidence-based interventions (EBI) in mental health lack sufficient evidence for specific racial and ethnic/cultural groups.

In recent years, however, there has been a growing literature evaluating the efficacy of various forms of psychotherapy and community-based interventions in diverse children and youth.^{71,84} It is incumbent on clinicians to preferentially utilize EBIs with specific population-based evidence. Psychotherapeutic interventions demonstrating evidence with diverse populations include cognitive behavioral therapy for treatment of depression in Latinos and African-Americans,^{72,85} interpersonal psychotherapy for treatment of depression in Latinos,⁸⁵ cognitive behavioral therapy for treatment of anxiety disorders in Latinos,⁸⁶ group cognitive behavioral therapy for anxiety disorders in African-Americans,⁸⁷ trauma-focused cognitive psychotherapy and peer-mediated treatment for traumatic stress for African-Americans,⁸⁸ and manualized family therapy for the treatment of substance abuse for Latinos.⁸⁹ Most of these studies have involved culturally adapted protocols. Community-based interventions with racial/ethnic specific evidence include Multi-Systemic Therapy for conduct disturbances,

substance abuse, and suicidality in African-Americans, Asian-Pacific Islanders and Latinos,⁹⁰ school-based cognitive behavioral interventions for treatment of depression in African-Americans and Latinos,⁹¹ and school-based cognitive behavioral interventions for trauma in Latinos.⁷⁵ A large number of EBIs have been evaluated with African American and Latino youth with conduct problems, with over a dozen of these distinct treatments having been successfully tested in randomized trials.⁷²

Relatively less evidence exists for pharmacological interventions; however, some data is available for diverse populations. For example, the Treatment of Adolescent Depression Study (TADS)⁹² had a 26% minority representation among its participants (principally Latino and African-American), and minority status was found not to be a significant moderator of acute treatment outcome using combined cognitive behavioral psychotherapy and pharmacotherapy. However, no separate data analyses on the effectiveness of the treatments examined have been published. In the Multisite Treatment Study of ADHD, inner city African-Americans and Latinos required combination stimulant pharmacotherapy and cognitive behavioral intervention to achieve equal outcomes to white children, who only required stimulant pharmacotherapy.⁹³

Principle 13. *Clinicians should identify ethnopharmacological factors (pharmacogenomic, dietary, use of herbal cures) that may influence the child's response to medications or their experience of side effects.*

It is important to avoid the harmful stereotyping around race and genetics of the past, but to adopt a new approach to the consideration of biological and genetic factors that serves the goal of addressing health disparities.⁹⁴ Although most genetic variation is shared worldwide, the relative proportion of functional genetic variants for any given gene may vary by ancestry. This in theory can lead to different patterns of medication metabolism and activity, and risk for side effects. The area of ethnopsychopharmacology has focused on the study of pharmacogenomic risk alleles that vary in frequency across different ethnic and racial populations, raising questions about the importance of these factors in prescribing. These include the distribution of rapid, slow and super-slow activity of CPY isoenzymes (especially CPY 2D6 and CPY 2D19) across different racial and ethnic populations, and polymorphisms of the serotonin 2A and dopamine D3 receptors related to antipsychotic and antidepressant treatment response.^{95,96} These differences have been associated with reports of African-Americans (and Caribbean Latinos with African

heritage) experiencing lesser response to serotonin reuptake inhibitors and more frequent extrapyramidal side effects with antipsychotics, and Asians having a higher percentage of slow metabolizers (SM) and often experiencing Western medicines as being “too strong” with many side effects.^{96,97}

Despite extensive pharmacogenomic research, there are few identified genetic variants with definitive clinical utility.⁹⁸ There are a vast number of genes involved in medication response and metabolism, particularly for drugs with central nervous system activity. Much of the focus to date has been on particularly the effects of genes on enzymes responsible for Phase I metabolism (e.g., cytochrome P-450 [CYP] enzymes), receptor regulation, and receptor transporters. Genetic variants within these classes of genes have been variably associated with differential treatment response and side effect profiles. However, the effect sizes are small, and the specific risk variants are often not consistently replicated in other studies.⁹⁸

Using race or ethnicity as a solitary marker for specific pharmacogenomic profiles raises a number of confounds.⁹⁹ The human genome is enormously complex, the predominance of genetic variation in the human genome is shared across all human populations, yet individuals are genetically unique.¹⁰⁰ Alleles that vary widely in frequency across different human populations often have no functional significance.¹⁰¹ Alleles with widely varying frequencies based on ancestry are also prone towards false positive findings in genetic association studies, a phenomenon known as population stratification. Additionally, racial intermixing in modern populations renders generalizations around genetics meaningless.

One meta-analytic review of ethnic differences in tolerability of psychotropic drugs did not find significant differences.¹⁰² However, to date there has been significant under-recruitment of diverse populations into randomized controlled studies, thus limiting the validity and generalizability of such results. Newer studies focusing on pharmacogenetic variables, particularly with adults, are much more careful around participant recruitment, phenotypic classification, and controlled designs with adequate numbers of diverse participants to control for population stratification bias.⁹⁴ For example, using an ethnically stratified analysis to control for population stratification, a study of the putative role of the dopamine D3 receptor in tardive dyskinesia found that patients homozygous for the glycine variant of the DRD3 gene had significantly higher AIMS scores, and African-Americans had a significantly higher prevalence of this polymorphism.¹⁰³

Gene-environment interactions are also important in predicting medication response, and further complicate the use of race or ethnicity as a simple pharmacogenomic predictor.⁹⁹ For example, grapefruit juice may increase serum concentrations of nefazodone and alprazolam by affecting the CYP3A4 isoenzymes and corn diets may increase serum concentrations of selective serotonin reuptake inhibitors by affecting CYP2D6 isoenzymes.⁹⁷ Thus, the dietary practices of different cultural groups need to be considered when prescribing.⁹⁶ Such practices may change as individuals migrate to different regions and adopt the practices of their new community. Additionally, findings of interactions between stressful life events and genetic polymorphisms in serotonin transporter genes in expression of major depression may present even greater confounds for minority and underserved populations chronically exposed to such stress.¹⁰⁴

Thus, at this time, although there is great promise for pharmacogenomics to advance personalized medicine, further research is needed to identify genetic factors with definitive impact on treatment response and side effect profiles. Until such findings are available, clinicians should exercise caution in prescribing and dosing psychopharmacological agents for diverse patients. They should base decisions on the individual's clinical and familial history, migration and ancestral history, personal and familial history of pharmacological response, and dietary history.⁹⁶ Currently available pharmacogenomic testing may have some limited utility in cases of diverse children with mixed racial/ethnic background and atypical pharmacological response.⁹⁸

It is important that clinicians inquire about and discuss the use of alternative medicinals and herbals by culturally diverse youth and families. Western medicine has not eclipsed traditional medicine, and the two are often practiced simultaneously or sequentially. Herbs and other traditional remedies sometimes have strong active ingredients, such as atropinic substances that can produce anti-cholinergic side effects and even toxicity. Many patients and families do not inform their doctor they are consuming these substances unless asked directly.⁹⁶ When these do not pose any adverse interaction or might even be beneficial, clinician acceptance of their use can help to build the therapeutic alliance and acceptance of Western therapeutic approaches.

PARAMETER LIMITATIONS

AACAP Practice Parameters are developed to assist clinicians in psychiatric decision-making. These parameters are not intended to define the sole standard of care. As such, the parameters should not be deemed inclusive of all proper methods of care nor exclusive of other

methods of care directed at obtaining the desired results. The ultimate judgment regarding the care of a particular patient must be made by the clinician in light of all of the circumstances presented by the patient and his or her family, the diagnostic and treatment options available, and available resources.

Disclosures: Dr. Pumariega has received research support from Eli Lilly and Company and has participated on an advisory panel on ADHD in Latino children from Shire Pharmaceuticals. Dr. Rothe has or had a leadership position with the American Academy of Psychoanalysis and Dynamic Psychiatry. Dr. Mian has or had a leadership position with the International Association of Child and Adolescent Psychiatry and Allied Professions. Dr. Toppelberg has received research support from the Norwegian Institute of Public Health, the Norwegian Research Council, and the National Institute of Mental Health; he also has participated on an advisory panel on ADHD in Latino children from Shire Pharmaceuticals. Dr. Harris has or had leadership roles with the Association for Academic Psychiatry, Black Psychiatrists of America, AAMC, and the American Psychiatric Association. Dr. Gogineni has or had a leadership position with the American Association for Social Psychiatry. Drs. Carlisle, Webb and Smith have no leadership or financial conflicts to disclose. Dr. Bukstein has or had a leadership role with the American Psychiatric Association, has intellectual property with Routledge Press, and serves or has served as an advisor/consultant to Ezra Innovations. Dr. Walter has no leadership or financial conflicts to disclose.

REFERENCES

1. U.S. Census (2009) United States Population Projections: 2000 to 2050. <http://www.census.gov/population/www/projections/analytical-document09.pdf>. Accessed June 22, 2011
2. Four Racial/Ethnic Panels. *Cultural Competence Standards for Managed Care Mental Health for Four Racial/Ethnic Underserved/Underrepresented Populations*. Rockville, MD: Center for Mental Health Services, Substance Abuse and Mental Health Administration, U.S. Department of Health and Human Services; 1999.
3. Cross T, Bazron B, Dennis K, Issac M. *Towards a Culturally Competent System of Care*. Washington, DC: CASSP Technical Assistance Center, Georgetown University Child Development Center; 1989.
4. Rothe EM, Tzuang D, Pumariega AJ. Acculturation, development and adaptation. *Child Adolesc Psychiatr Clin N Am*. 2010;19:681-696.
5. Eaton D, Kann L, Kinchen S, et al. Youth risk behavior surveillance-United States 2009. *Morbidity and Mortality Weekly Report Surveillance Summaries* Vol. 59 / No. SS-5, 1-142. Published June 4, 2010. <http://www.cdc.gov/mmwr/pdf/ss/ss5905.pdf> . Accessed June 22, 2011
6. Institute of Medicine. *Unequal Treatment: Confronting Racial and Ethnic Disparities in Health Care*. Washington: National Academies Press; 2002.
7. U.S. Office of the Surgeon General. *Mental health: Culture, race, and ethnicity. A supplement to: Mental health: A report of the surgeon general*. Rockville, MD: U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, National Institutes of Health, National Institute of Mental Health; 2001.
8. Alegria M, Vallas M, Pumariega A. Racial and ethnic disparities in pediatric mental health. *Child Adol Psych Clin*. 2010;19(4):759-774.
9. Angold A, Erkanli A, Farmer E, et al. Psychiatric disorder, impairment, and service use in rural African-American and white youth. *Arch Gen Psychiatry*. 2002;59(10):893-901.
10. Cuffe S, Waller J, Cuccaro M, Pumariega A. Race and gender differences in the treatment of psychiatric disorders in young adolescents. *J Am Acad Child Adolesc Psychiatry*. 1995;34(11):1536-43.
11. Pumariega A, Glover S, Holzer C, Nguyen N. Utilization of mental health services in a tri-ethnic sample of adolescents. *Comm Mental Health J*. 1998;34(2):145-156.
12. Costello E, Pescosolido B, Angold A, Burns BJ. A family network-based model of access to child mental health services. *Research in Community and Mental Health*. 1998;9:165-190.
13. Snowden L, Yamada A. Cultural differences in access to care. *Annu Rev Clin Psychol*. 2005;1:143-66.
14. Kouyoumdjian H, Zamboanga B, Hansen D. Barriers to community mental health services for Latinos: Treatment considerations. *Clinl Psychol- Sci Pr*. 2003;10(4):394-422.
15. Abe-Kim J, Takeuchi D, Hong S, et al. Use of mental health-related services among immigrant and US-born Asian Americans: Results from the national Latino and Asian American study. *Am J Public Health*. 2007;97(1):1-8.
16. Freedenthan S, Stiffman A. "They might think I was crazy": Young American Indians' reasons for not seeking help when suicidal. *J Adol Research*. 2007; 22:58-77.

17. Suite D, LaBril R, Primm A, Harrison-Ross P. Beyond misdiagnosis, misunderstanding and mistrust: Relevance of the historical perspective in the medical and mental health treatment of people of color. *J Natl Med Assoc.* 2007; 99:879-885.
18. Napoles-Springer A, Santavo J, Houston K, Perez-Stable E, Stewart A. Patients' perceptions of cultural factors affecting the quality of their medical encounters. *Health Expect.* 2005;8(1):4-17.
19. Yeh M, McCabe K, Hurlburt M, et al. Referral sources, diagnoses, and service types of youth in public outpatient mental health care: a focus on ethnic minorities. *J Behav Health Ser R.* 2002;29:45–60.
20. Sentell T, Shumway M, Snowden L. Access to mental health treatment by English language proficiency and race/ethnicity. *J Gen Intern Med.* 2007;22(2):289–293
21. Malgady R, Constantino G. Symptom severity in bilingual Hispanics as a function of clinician ethnicity and language of interview. *Psychol Assessment.* 1998;10(2):120-127.
22. Flores G, Laws M, Mayo S, et al. Errors in medical interpretation and their potential clinical consequence in pediatric encounters. *Pediatrics.* 2003;111:6–14.
23. Martinez C, McClure H, Eddy, J. Language brokering contexts and behavioral and emotional adjustment among Latino parents and adolescents. *J Early Adol.* 2009;29:71-98.
24. Pumariega A, Rothe E, Rogers K. Cultural competence in child psychiatric practice. *J Am Acad Child and Adolesc Psychiatry.* 2009;48(4):362-366.
25. Toppelberg C, Munir K, Nieto-Castañon A. Spanish-English bilingual children with psychopathology: Language deficits and academic language proficiency. *Child and Adolescent Mental Health.* 2006; 11:156-163.
26. Toppelberg C, Collins B. Language, culture and adaptation in immigrant children. *Child Adol Psych Cl.* 2010;19(4):697-717.
27. Toppelberg C, Tabors P, Coggins A, Lum K, Burger C. Differential diagnosis of selective mutism in bilingual children. *J Am Acad Child Adolesc Psychiatry.* 2005;44(6):592-595.
28. Kilgus M, Pumariega A, Cuffe S. Influence of race on diagnosis in adolescent psychiatric patients. *J Am Acad Child Adolesc Psychiatry.* 1995;34(1):67-72.
29. Van Ryn M. Research on the provider contribution to race/ethnicity disparities in medical care. *Medical Care.* 2002;40(1):140-151.
30. Sue D, Capodilupo C, Torino G, Bucceri J, Holder A, Nadal K, Esquilin M. Racial microaggressions in everyday life: Implications for clinical practice. *Am Psychol.* 2007;62(4): 271–286.
31. Dovidio J, Brigham J, Johnson B, Gaertner S. Stereotyping, prejudice, and discrimination: Another look. In Macrae N, Stangor C, Hewstone M eds., *Stereotypes and Stereotyping.* New York, NY: Guilford Press; 1996: 276-319.
32. Abreu J. Conscious and nonconscious African American stereotypes: Impact on first impression and diagnostic ratings by therapists. *J Consult Clinl Psych.* 1999;67(3):387-93.
33. Pumariega A, Joshi S. Culture and development. *Child Adol Psych Clin.* 2010;19(4): 661-680.
34. Kirmayer L. Cultural variations in the clinical presentation of depression and anxiety: implications for diagnosis and treatment. *J Clin Psychiatry* 2001;62 [suppl.13]:22–28.
35. Malgady R, Rogler L, Dharma E. Cultural expression of psychiatric symptoms: Idioms of anger amongst Puerto Ricans. *Psychol Assmt.* 1996; 8(3): 265-268.

36. Chentsova-Dutton Y, Tsai J, Chu J, Rottenberg J, Gross J, Gotlib I. Depression and emotional reactivity: Variation among Asian Americans of East Asian descent and European-Americans. *J Abnorm. Psychol.* 2007; 116:776-785.
37. American Psychiatric Association. Diagnostic and statistical manual of mental disorders, 4th edition, text revision (DSM IV-TR). Washington: American Psychiatric Association; 2000: 897-904.
38. Aggarwal, N. Cultural formulations in child and adolescent psychiatry. *J Am Acad Child Adolesc Psychiatry.* 2010;49:306-309.
39. American Psychiatric Association. Diagnostic and statistical manual of mental disorders, 5th edition (DSM 5). Washington: American Psychiatric Association; 2013. In Press.
40. American Academy of Child and Adolescent Psychiatry. Practice parameter on gay, lesbian, or bisexual sexual orientation, gender nonconformity, and gender discordance in children and adolescents. *J Am Acad Child Adolesc Psychiatry.*2012; 51: 956-974.
41. Sack W. Multiple forms of stress in refugee and immigrant children. *Chil Adol Psych Clin.* 1998;7:153-167.
42. Orfield G, Yun J. Resegregation in American schools. Cambridge, MA: Civil Rights Project, Harvard University; 1998.
43. Rothe EM, Lewis J, Castillo-Matos H, Martinez O, Busquets R, Martinez I. Post traumatic stress disorder in Cuban children and adolescents after release from a refugee camp. *Psychiatric Services.* 2002;53:970-976.
44. Pumariega AJ, Rothe EM. Leaving no children or families outside: the challenges of immigration. *Am J Orthopsychiatry* 2010;80:506-516.
45. Suarez-Orozco C, Suarez-Orozco M. *Children and Immigration.* Cambridge, MA: Harvard University Press; 2001.
46. Vigil D. Barrio Gangs: Street Life and Identity in Southern California. Austin, TX: University of Texas Press; 1988.
47. Hovey J, King C. Acculturative stress, depression, and suicidal ideation among immigrant and second-generation Latino adolescents. *J Am Acad Child Adolesc Psychiatry.* 1996;35:1183-1192.
48. McFarlane J, Groff J, O'Brien J, Watson K. Behaviors of children who are exposed and not exposed to intimate partner violence: an analysis of 330 Black, White and Hispanic children. *Pediatrics.* 2003;112:202-207.
49. Suarez-Orozco C, Todorova I, Louie J. (2002) Making up for lost time: the experience of separation and reunification among immigrant families. *Fam Process.* 2002;41(4):625-643.
50. Almqvist K, Broberg A. Mental health and social adjustment in young refugee children 3 1/2 years after their arrival in Sweden. *J Am Acad Child Adolesc Psychiatry.* 1999;38(6):723-730.
51. Lustig SL, Kia-Keating M, Grant-Knight W, et al. Review of child and adolescent psychiatry refugee mental health. *J Am Acad Child Adolesc Psychiatry.* 2004;43:24-36.
52. Erikson EH. (1968). *Identity: Youth and Crisis.* New York: Norton.
53. Duarte CS, Bird HR, Shrout PE, Wu P, Lewis-Fernandéz R, Shen S, Canino G. Culture and psychiatric symptoms in Puerto Rican children: longitudinal results from one ethnic group in two contexts. *J Child Psychol Psychiatry.* 2008;49:563-72.
54. Becker DF, Weine SM, Vojvoda D. PTSD symptoms in adolescent survivors of "ethnic cleansing": results from a one-year follow-up study. *J Am Acad Child Adolesc Psychiatry.* 1999;38:775-781.

55. Szapocznik J, Kurtines W. *Breakthroughs in Family Therapy with Drug Abusing and Problem Youth*. New York: Springer, 1989.
56. Sullivan S, Schwartz S, Prado G, Huang S, Pantin H, Szapocznik JA. Bidimensional model of acculturation for examining differences in family functioning and behavior problems in Hispanic immigrant adolescents. *J. Early Adolesc.* 2007;27:405-430.
57. Hwang, W., Wood, J. Acculturative family distancing: Links with self-reported symptomatology among Asian Americans and Latinos. *Child Psychiat Human Behav.* 2009;40:123–138.
58. Chatters, L.M., Taylor, R.J., Jayakody, J.S. Fictive kinship relationships in Black extended families. *J Comp Fam. Stud.* 1994;25:297-312.
59. Abe-Kim J, Takeuchi D, Hwang W. Predictors of help seeking for emotional distress among Chinese Americans: family matters. *J Consult Psychol.* 2002;70:1186–90.
60. Myers, GE. Addressing the effects of culture on the boundary-keeping practices of psychiatry residents educated outside of the United States. *Acad. Psychiatry.* 2004; 28:47-55.
61. Savin D, Martinez R. Cross-cultural boundary dilemmas: a graded-risk assessment approach. *Transcult. Psychiatry,* 2006;43:243-258.
62. Snowden LR. Explaining mental health treatment disparities: ethnic and cultural differences in family involvement. *Cult. Med. Psychiatry.* 2007;31: 389-402.
63. Sue S. In defense of cultural competency in psychotherapy and treatment. *Am Psychol.* 2003;58: 964–970.
64. Ngo-Metzger Q, Telfair J, Sorkin DH, et al. (2006) *Cultural Competency and Quality of Care: Obtaining the Patient's Perspective*. The Commonwealth Fund. Publication No. 963, p. 7-9. www.cmwf.org, last accessed on 9/6/2011.
65. Costantino G, Malgady RG, Rogler LH. Storytelling through pictures: Culturally sensitive psychotherapy for Hispanic children and adolescents. *J Clin Child Psychol.* 1994;23:13–20.
66. Jackson-Gilfort A, Liddle HA, Tejeda MJ, Dakof GA. Facilitating engagement of African American male adolescents in family therapy: A cultural theme process study. *J Black Psychol.* 2001;27:321–340.
67. Moran JR. Preventing alcohol use among urban American Indian youth: The seventh generation program. *J Hum Behav Soc Environ.* 1999;2:51–67.
68. Harvey AR, Hill RB. Afrocentric youth and family rites of passage program: Promoting resilience among at-risk African American youths. *Soc Work,* 2004;49:65–74.
69. Halliday-Boykins CA, Schoenwald SK, Letourneau EJ. Caregiver-therapist ethnic similarity predicts youth outcomes from an empirically based treatment. *J Consult Clin Psychol.* 2005;73:808–818.
70. Yeh M, Eastman K, Cheung MK. Children and adolescents in community health centers: Does the ethnicity or the language of the therapist matter? *J Comm Psychol.* 1994;22:153–163.
71. Griner D, Smith T. Culturally adapted mental health interventions: a meta-analytic review. *Psychotherapy: Theory Res Pract Train.* 2006;43:531–548.
72. Huey S, Polo A. Evidence-based psychosocial treatments for ethnic minority youth. *J Clin Child Adol Psychol.* 2008;37:262–301.
73. Akutsu PD, Castillo ED, Snowden LR. Differential referral patterns to ethnic-specific and mainstream mental health programs for four Asian American groups. *Am J Orthopsychiatry.* 2007;77:95-103.

74. Dean KL, Jaycox LH, Langley AK, Kataoka SH, Stein BD. School-based disaster mental health services: Clinical, policy and community challenges. *Professional Psychology: Research and Practice*. 2008;39:52-57.
75. Kataoka S, Stein B, Jaycox L, et al. A school-based mental health program for traumatized Latino immigrant children. *J Am Acad Child Adolesc Psychiatry*. 2003;42:311-318.
76. American Academy of Child and Adolescent Psychiatry. Practice parameters for child mental health in systems of care. *J Am Acad Child Adol Psychiatry*. 2007;46:284-299.
77. Miech R, Azur M, Dusablon T, et al. The potential to reduce mental health disparities through the comprehensive community mental health services for children and their families program. *J Behav Health Serv Res*. 2008;35:253-264.
78. Fisher S, Sukumar B, Manteuffel B, Stephens R. A preliminary profile of Latino children and youth receiving services in systems of care communities. *Psychline*. 2005;4:4-12.
79. Martinez CR, Eddy JM. Effects of culturally adapted parent management training on Latino youth behavioral health outcomes. *J Couns Clin Psychol*. 2005;73:841-851.
80. Kumpfer KL, Alvarado R, Smith P, Bellamy N. Cultural sensitivity and adaptation in family-based prevention interventions. *Prevent Science* 2002;3:241-246.
81. Jambunathan S, Conselman KP. Parenting attitudes of Asian Indian mothers living in the United States and in India. *Early Child Develop Care*. 2002;172:657-662.
82. Ali SR, Liu WM, Humedian M. Islam 101: Understanding the religion and therapy implications. *Prof Psychol Res Pr*. 2004;35:635-642.
83. Rothbaum F, Morelli G, Pott M, Liu-Constant Y. Immigrant-Chinese and Euro-American parents' physical closeness with young children: Themes of family relatedness. *J Fam Psychol*. 2000;14:334-348.
84. Kataoka S, Novins D. The practice of evidence-based treatments in ethnic minority youth. *Child Adol Psychiatr Clin N Am*. 2010;19:775-790.
85. Pumariega AJ, Rothe EM, Rogers KM. Depression in immigrant and minority children. In: J M Rey & J Birmaher (Eds.), *Treating Childhood Depression*. New York: Lippincott, Williams and Wilkins, 2008.
86. Pina A, Wendy M, Silverman K, Fuentes R, Kurtines W, Weems C. Exposure-based cognitive-behavioral treatment for phobic and anxiety disorders: Treatment effects and maintenance for Hispanic/Latino relative to European-American youths. *J Am Acad Child Adolesc Psychiatry*. 2003;42:1179-1187.
87. Ginsburg GS, Drake KL. School-based treatment for anxious African Americans adolescents: A controlled pilot study. *J Am Acad Child Adolesc Psychiatry*. 2002;41:768-775.
88. Cohen JA, Deblinger E, Mannarino AP, Steer RA. A multisite, randomized controlled trial for children with sexual abuse-related PTSD symptoms. *J Am Acad Child Adolesc Psychiatry*. 2004;43:393-402.
89. Santisteban DA, Coatsworth JD, Perez-Vidal A, et al. Efficacy of brief strategic family therapy in modifying Hispanic adolescent behavior problems and substance use. *J Fam Psychol*. 2003;17:121-133.
90. Rowland MD, Halliday-Boykins CA, Henggeler SW, et al. A randomized trial of multisystemic therapy with Hawaii's Felix Class youths. *J Emot Behav Disord*. 2005;13:13-23.
91. Cardemil E, Reivich K, Beevers C, Seligman M, James J. The prevention of depressive symptoms in low-income minority children: Two year follow-up. *Behav Res Ther*. 2007;45:313-327.

92. Curry J, Rohe P, Simons A, et al. Predictors and modifiers of acute outcome in the Treatment for Adolescents with Depression Study (TADS). *J Am Acad Child Adolesc Psychiatry*. 2006;45:1427-1439.
93. Arnold LE, Elliott M, Sachs L, et al. Effects of ethnicity on treatment attendance, stimulant response-dose, and 14-month outcome in ADHD. *J Consult Clin Psychol*. 2003;71:713-727.
94. Shields A, Fortun M, Hammonds E, et al. The use of race variables in genetic studies of complex traits and the goal of reducing health disparities: A transdisciplinary perspective. *Am Psychol*. 2005;60:77-103.
95. McMahon F, Buervenich B, Charney D, et al. Variation in the gene encoding the serotonin 2A receptor is associated with outcome of antidepressant treatment. *Am J Hum Genet*. 2006;78:804-814.
96. Malik M, Lawson W, Lake J, Joshi S. Culturally adapted pharmacotherapy and the integrated formulation. *Child Adolesc Psychiatr Clin N Am*. 2010;19:791-814.
97. Lin K, Gray G. Ethnopharmacology in Asians. In: Pedro Ruiz, ed. *Ethnicity and Psychopharmacology*. Washington, D.C.: American Psychiatric Press, 2000.
98. Zandi PP, Judy JT. The promise and reality of pharmacogenetics in psychiatry. *Clin Lab Med*. 2010;30(4):931-74.
99. Jones DS, Perlis RH. Pharmacogenetics, race, and psychiatry: Prospects and challenges. *Harv Rev Psychiat*. 2006;14(2): 92-108.
100. Cavalli-Sforza LL, Menozzi P, Piazza A. *History and Geography of Human Genes*. Princeton University Press, Princeton, NJ, 1994.
101. McClellan J, King MC. Genetic heterogeneity in human disease. *Cell*. 2010;141(2):210-7.
102. Ormerod S, McDowell SE, Coleman JJ, Ferner RE. Ethnic differences in the risks of adverse reactions to drugs used in the treatment of psychoses and depression: a systematic review and meta-analysis. *Drug Saf*. 2008;31(7):597-607.
103. Basile V, Masellis M, Badri F, et al. Association of the *MscI* polymorphism of the dopamine D3 receptor gene with tardive dyskinesia in schizophrenia. *Neuropsychopharm*. 1999;21:17-27.
104. Kendler K, Kuhn J, Vittum J, Prescott C, Riley B. The interaction of stressful life events and a serotonin transporter polymorphism in the prediction of episodes of major depression: a replication. *Arch. Gen. Psychiatry*. 2005;62:529-535.